

COBBER G



Organic Copper
Foliar / Soil



Fungicide - Bactericide
Cu Deficiency Corrector

CHARACTERISTICS

COBBER G contains the copper ion (Cu^{+2}) complexed by a natural organic molecule (Gluconic acid). This complexation favors the transport of Cu in the plant.

COBBER G is a systemic product that stimulates the synthesis of PHYTOALEXINES, which increase the natural defenses of the crops.

COMPOSITION

	%w/w	%w/v
Copper (Cu)	6,5	8
Organic Complexant Agent Gluconic Acid		

- **BIODEGRADABLE**
- **SYSTEMIC ACTION**
- **DOES NOT STAIN THE PLANT**
- **IMPROVE CONSERVATION OF FRUITS**
- **HIGH ASSIMILATION**

ACTIONS

PREVENTS THE ABIOTIC STRESS PLANT

FUNGICIDE / NATURAL ORGANIC BACTERICIDAL

HIGH EFFICIENCY

APPLICATION

Crops	Application
Cereals	Foliar: 150-250 cc/hL
	Foliar: 200-300 cc/hL (2 to 3 application from the beginning of budding)
Citrics	Fertirrigation: 6L-12L/ha (Apply 2-3 times: the first application near vegetative cessation, the second at the onset of budding, and the last during full vegetative activity)
	Foliar: 200-300 cc/hL (2 to 3 application from the beginning of budding)
Fruit Trees	Fertirrigation: 6L-12L/ha (Apply 2-3 times: the first application near vegetative cessation, the second at the onset of budding, and the last during full vegetative activity)
	Foliar: 150-300 cc/hL (3-4 applications as needed and based on the crop's development)
Horticultural Crops	Fertirrigation: 5L-10L/ha (3-4 applications as needed and based on the crop's development)
	Foliar: 200-300 cc/hL (2 to 3 application from the beginning of budding)
Olive Trees	Fertirrigation: 6L-12L/ha (Apply 2-3 times: the first application near vegetative cessation, the second at the onset of budding, and the last during full vegetative activity)
Rice	Foliar: 150-250 cc/hL Fertirrigation: 5L/ha Copper deficiency
Vine	Foliar: 200-300 cc/hL until veraison

Cautions

It is recommended to treat between 6 and 25 °C. Avoid applications in cases of extreme drought, humidity, frost and rain. Shake well before use. Do not freeze. In case of mixing with other products, always carry out a previous test. Incorporate this product into the last phase.

