



**Aminoacid
Foliar/Solid**



SOWY

COMPOSITION

	%w/w
Total Aminoacids	28
Free Aminoacids	25
Total Nitrogen	5
Organic Nitrogen	4
Defense inductor (ID)	3

- **VEGETAL ORIGIN**
- **BIOSTIMULATING ACTIVITY**
- **ANTISTRESS EFFECT**

Liquid fertilizer of high concentration of amino acids

CHARACTERISTICS

SOWY is a product of natural origin, with a high content of free amino acids, derived from vegetable matter. Composed of amino acids that the plant uses to satisfy its protein needs, with an important energysaving, acting as an effective biostimulant during the rooting, sprouting, flowering, fruit setting and fruit development stages.

APPLICATION

CROPS	SOIL DOSAGE	Lts/ha
Strawberries	Every 10 days after transplanting	4
Fruit trees	From budding until the swelling of the fruit	6
Banana plants	Every 15 days between March and June	6
Olive trees	Throughout the whole cycle	18
Table grapes	From budding until the end of the cycle	5
Dry fruits	From budding until the swelling of the fruit	5
Citrus fruit	From flowering until the swelling of the fruit	12
Cotton	10 days after shooting until 20 days after the flowering	6
Ornamental Plants	Every 15 days after transplanting	4

ACTIONS

- ✓ **INCREASED PRODUCTIVITY**
- ✓ **FAVORIZES RADICULAR DEVELOPMENT**
- ✓ **IMPROVES THE ABSORPTION AND TRANSPORTATION OF NUTRIENTS**
- ✓ **INDUCES A BETTER POLLINATION AND FLOWERING**
- ✓ **GOOD SPROUTING**
- ✓ **FRUITS OF GREATER QUALITY**

CROPS	FOLIAR DOSAGE	cc/100L
Horticultural crops	Every 10 days after transplanting	200
Strawberries	Throughout the whole cycle	200
Tubers	Every 15 days	250
Fruit trees	From budding until the swelling of the fruit	200-300
Banana plants	Every 15 days	250
Olive trees	Throughout the whole cycle	200-300
Table grapes	From budding until the end of the cycle	250
Wine grapes	From budding until the end of the cycle	2 L/Ha
Dry fruits	From budding until the swelling of the fruit	200-300
Citrus fruits	From flowering until the swelling of the fruit	200-300
Beet	2 applications every 15 days	2,5 L/Ha
Cotton	10 days after sprouting until 20 days after the first flower.	300
Alfalfa	After every mowing	2,5 L/Ha
Ornamental plants	Every 15 days after transplanting	250
Lawn	After sowing/Growth phase	3-5 L/Ha/ 30 cc/m ²

