tangelagro

Datasheet



SOWY Solid

COMPOSITION	%w/w	
Free aminoacids	82,00	
Total Nitrogen (N)	12,00	

- SOLID OF HIGH SOLUBILITY
- VEGETAL ORIGIN
- BIOSTIMULATING ACTIVITY
- ANTISTRESS EFFECT

Solid free amino acids

CHARACTERISTICS

SOWY SOLID is a biostimulant product composed of amino acids of vegetable origin obtained by enzymatic hydrolysis of natural proteins. Stimulates the essential metabolic processes in a crop: rooting, sprouting, flowering, setting and fruit development, especially when there are adverse situations in the process of plant development.

ACTIONS

- INCREASE RESISTANCE TO STRESS (High temperature, low humidity, drought, pest attacks, frost, flood)
- **✓** PROTEIN SYNTHESIS
- INCREASES CONTENT OF CHLOROPHYLL
- **REGULATES STOMATAL OPENING**
- CHELATING AGENT FOR MICRONUTRIENTS
- RAW MATERIALS HORMONES
- HELPS POLLINATION AND FRUIT SET
- INCREASES SOIL MICROBIAL ACTIVITY

SOIL APPLICATION

CROPS	Application	CROPS	Application
In all crops	SEASON: Reduction of stress, improvement of efficiency of plant protection treatment ANNUAL DOSAGE: 3-4 Kg/Ha in case of stress (salinity, dryness, etc) divided into several doses (1 Kg/Ha)	Open field vegetables	SEASON: Yield increase ANNUAL DOSAGE: 1-2 Kg/Ha every 7-10 days after the first true leaf stage
		Ornamental plants and tree nursery, landscaping, turf	SEASON: Root formation and nutrient uptake, sprouting leaf quality, growth ANNUAL DOSAGE: 1 Kg/Ha every 7-12 days starting from planting SEASON: Yield, sprouting leaf quality, growth ANNUAL DOSAGE: 3-4 Kg/Ha divided into 2-3 applications every 10-15 days, starting from planting
Cereals, potatoes, legumes	SEASON: Yield increase ANNUAL DOSAGE: 3-4 Kg/Ha divided into 3 applications every 15 days after the first true leaf stage	grass (in general) Vegetables in greenhouses	
Horticultural fruit trees	SEASON: Fruit setting, fruit growth and quality ANNUAL DOSAGE: 1-2 Kg/Ha every 10-15 days, from pre blooming stage until the beginning of the colouring stage		

The recommended concentration for **foliar application** is 0,3-0,5% in the usual quantity of spray water.



