

Datasheet



Inorganic Copper
Foliar / Soil



Multisite Contact Activity
MCA

BioFungicide - Bactericide
Cu Deficiency Corrector

COBBER S

COMPOSITION

%w/v

Copper (Cu)	20 (200 g/L)
Copper sulfate	75 (750 g/L)
Sulfur (SO ₃)	26 (260 g/L)
Density:	1,4
pH:	4,5 - 5

- **MULTISITE CONTACT ACTIVITY**
- **EXCELLENT CROP SAFETY AND LOW USE RATES**
- **CONSISTENT, SUB-MICRON PARTICLE SIZE FOR BETTER COVERAGE AND DISEASES CONTROL**
- **BETTER ADHESION AND COVERAGE**

CHARACTERISTICS

COBBER S is a fungicide and contact bactericide, broad spectrum, preventive and curative action for the control of some diseases.

The smaller particle size in **COBBER S** delivers better plant coverage, which means better protection against fungal and bacterial diseases. The premium formulation readily mixes in water and stays suspended longer than any other liquid formulation.

ACTIONS: KEY DISEASES CONTROLLED

 <i>Alternaria</i>	<i>Collectrichum spp.</i>	<i>Phomopsis stem blight</i>
 <i>Anthracnose</i>	<i>Dieback</i>	<i>Phytophthora bud rot</i>
 <i>Bacterial leaf spot</i>	<i>Downy mildew</i>	<i>Pseudomonas leaf spot</i>
 <i>Bacterial soft rot</i>	<i>Entomosporium</i>	<i>Scab</i>
 <i>Bacterial spot</i>	<i>Exosporium leaf spot</i>	<i>Septoria leaf spot</i>
 <i>Black spots</i>	<i>Fire blight</i>	<i>Volutella leaf blight</i>
 <i>Botrytis</i>	<i>Leaf spot</i>	<i>Xanthomonas leaf spot</i>
 <i>Cercospora leaf spot</i>	<i>Pestalotia leaf spot</i>	

FOLIAR APPLICATION

Crops	Dosage
Citrus	75-125 cc/HL
Fruits Trees (Winter)	250-400 cc/HL
Forest nurseries	150-180 cc/HL
Herbaceous and Ligneous	150-250 cc/HL

Crops	Dosage
Olive	300-600 cc/HL
Pistachio	200-400 cc/HL
Vegetable	150-180 cc/HL
Vine	200-300 cc/HL

Caution

Compatible with most insecticides and fungicides. Do not mix with acides or alkalis. Do not add amino acids. Shake well before use. Do not freeze.

