



Nutrient
corrector
Soil / Foliar



TANGEL MIX MICRO

COMPOSITION

%w/w

Iron (Fe)	7,50
Manganese (Mn)	3,50
Zinc (Zn)	0,70
Boron (B)	0,65
Molybdenum (Mo)	0,30
Copper (Cu)	0,28
Chelating agent EDTA	

- TOTAL SOLUBILITY
- QUICK AND EFFECTIVE ASSIMILATION
- FOR ALL KIND OF CROPS

ACTIONS

- ✓ CORRECTS SEVERES MICRONUTRIENTS DEFICIENCIES. YIELD AND QUALITY IN CROPS.
- ✓ EDTA (CHELATING AGENT), FACILITATES THE UPTAKE AND TRANSPORT TO THE PLANT.

Solid fertilizer corrector of multiple deficiencies

CHARACTERISTICS

TANGEL MIX MICRO is a solid compound, highly-soluble in all types of water and whose Iron, Manganese, Copper, Zinc, Boron, Molybdenum and Magnesium micronutrients contribute simultaneously to the plant by providing the necessary dosage of nutrients that are indispensable for the perfect development of any crop.

Except for the Boron and Molybdenum, the nutrients in **TANGEL MIX MICRO** are included in a molecule (EDTA, ethylenediaminetetraacetic acid) that protects them in the soil and, when applied to the leaves, facilitates their uptake and transport to the plant.

APPLICATIONS

CROPS	Application	CROPS	Application
Cereals, Grain Legumes, Oilseed crops, Cotton, Maize, Summer crops	RATE: 1-1.5 kg/ha APPLICATION: Foliar CRITICAL COMMENTS: Apply early season, tillering and bolting.	Apples, Pears, Nut crops, Citrus, Mangoes, Stonefruit, Avocados, Pineapples.	APPLICATION: Foliar CRITICAL COMMENTS: Apply 2-3 times during the crop cycle. Apply by dilute application only. Do not exceed maximum per hectare rate. Minimum water rate of 500L per ha. RATE: 2-3 kg/ha APPLICATION: Fertigation CRITICAL COMMENTS: Apply every 7-10 days during the crop cycle.
Pastures, lucerne and forage crops.	RATE: 1-2 kg/ha APPLICATION: Foliar CRITICAL COMMENTS: Apply after each grazing or cutting.	Olives.	RATE: 0.5-1.5 kg/ha APPLICATION: Foliar CRITICAL COMMENTS: Apply when sufficient foliage is present. Apply in a minimum of 500L/ha of water. RATE: 5-10 kg/ha APPLICATION: Soil CRITICAL COMMENTS: Apply before sowing, transplanting or beginning of plant growth.
Potatoes and other tuber crops.	RATE: 1 kg/ha APPLICATION: Foliar CRITICAL COMMENTS: Apply up to 4 applications from 10cm stage until post flowering, at 7-10 day intervals. Apply in a minimum of 500L/ha of water.	Strawberries.	RATE: 2-3 kg/ha APPLICATION: Fertigation CRITICAL COMMENTS: Apply every 7-10 days during the crop cycle.
Vegetables	CRITICAL COMMENTS: Apply up to 3 applications at 10-15 day intervals when sufficient foliage is present. Apply in a minimum of 500L/ha of water. RATE: 5-10 kg/ha APPLICATION: Soil CRITICAL COMMENTS: Apply before sowing, transplanting or beginning of plant growth. RATE: 2-3 kg/ha APPLICATION: Fertigation CRITICAL COMMENTS: Apply every 7-10 days during the crop cycle. RATE: 2-3 kg/1000L APPLICATION: Hydroponics CRITICAL COMMENTS: Apply as required- 1L of prepared solution per 100 L irrigation water. Use the higher rate during crop development or periods of high temperature or during shorter and cooler days.	Other berry crops	RATE: 2-3 kg/1000L APPLICATION: Hydroponics CRITICAL COMMENTS: Apply as required- 1L of prepared solution per 100 L irrigation water. Use the higher rate during crop development or periods of high temperature or during shorter and cooler days.
Grapevines (wine and table).	APPLICATION: Foliar CRITICAL COMMENTS: Apply 2-3 times during the crop cycle. Apply by dilute application only. Do not exceed maximum per hectare rate. Minimum water rate of 500L per ha. RATE: 2-3 kg/ha APPLICATION: Fertigation CRITICAL COMMENTS: Apply every 7-10 days during the crop cycle.		

